

Supersymmetry of the magnetic vortex

C. DUVAL ⁽¹⁾ and P. A. HORVÁTHY ⁽²⁾

Abstract. *The $N = 2$ supersymmetry of the Pauli Hamiltonian in any static magnetic field in the plane combines, for the magnetic vortex, with Jackiw's bosonic $\mathfrak{o}(2) \times \mathfrak{o}(2,1)$ symmetry, into an $\mathfrak{o}(2) \times \mathfrak{osp}(1/2)$ dynamical supersymmetry.*

May 1993. Tours Preprint no 60/93.

A few years ago, Jackiw [1] pointed out that a spin-0 particle in a Dirac monopole field has an $\mathfrak{o}(2,1)$ dynamical symmetry, generated by the spin-0 Hamiltonian, $H_0 = \pi^2/2m$ where $\pi = \mathbf{p} - e\mathbf{A}$, by the dilatation and by the expansion,

$$(1) \quad D = tH_0 - \frac{1}{4}\{\pi, \mathbf{r}\} \quad \text{and} \quad K = -t^2H_0 + 2tD + \frac{m}{2}\mathbf{r}^2,$$

to which angular momentum adds an $\mathfrak{o}(3)$. This allowed him to calculate the spectrum and the wave functions group-theoretically [1].

Jackiw's result was extended to spin- $\frac{1}{2}$ particles by D'Hoker and Vinet [2] who have shown that, for the Pauli Hamiltonian $H = (1/2m) [\pi^2 - e\mathbf{B} \cdot \boldsymbol{\sigma}]$, not only the conformal generators D and K , but also the fermionic generators $Q = 1/\sqrt{2m} \pi \cdot \boldsymbol{\sigma}$ and $S = \sqrt{m/2} \mathbf{r} \cdot \boldsymbol{\sigma} - tQ$ are conserved. Thus, the spin system admits an $\mathfrak{o}(3) \times \mathfrak{osp}(1/1)$ conformal supersymmetry, yielding now an algebraic solution of the Pauli equation [2].

More recently, Jackiw [3] found that the $\mathfrak{o}(2,1)$ symmetry, generated by — formally — the same D and K as above, is also present for a magnetic vortex (an idealization for the Aharonov-Bohm experiment [4]), allowing for a group-theoretic treatment of the problem.

In this Letter we show that the $N = 2$ supersymmetry of the Pauli Hamiltonian of a spin- $\frac{1}{2}$ particle (present for any magnetic field in the plane [5]) combines, for a magnetic vortex, with Jackiw's $\mathfrak{o}(2) \times \mathfrak{o}(2,1)$ into an $\mathfrak{o}(2) \times \mathfrak{osp}(1/2)$ superalgebra. This result is to be compared with the Galilean supersymmetry discovered recently by Leblanc et al. [6] for non-relativistic Chern-Simons systems, and with the $\mathfrak{osp}(1/2)$ found by Hughes et al. for a constant magnetic field [7].

Thus the planar system has more symmetries as its higher-dimensional counterpart. We call this supersymmetry exotic, because it is realized with two — rather than four-component — objects. Our pseudo-classical calculations in Ref. [6] indicate that such an 'exotic' supersymmetry is only possible in two spatial dimensions — one more indication of the particular status of two-dimensional physics.

Let us start with a spin- $\frac{1}{2}$ particle in a static magnetic field $\mathbf{B} = (0, 0, B(x, y))$. Dropping the irrelevant

⁽¹⁾ Centre de Physique Théorique, CNRS, Luminy, Case 907. F-13288 Marseille Cedex 9 (France). UMR 6207 du CNRS associée aux Universités d'Aix-Marseille I et II et Université du Sud Toulon-Var; Laboratoire affilié à la FRUMAM-FR2291. mailto: duval@cpt.univ-mrs.fr.

⁽²⁾ Laboratoire de Mathématiques et de Physique Théorique, Université de Tours, Parc de Grandmont, F-37200 Tours (Fr). mailto: horvathy@lmpt.univ-tours.fr.

z variable, we can work in the plane. Then our model is described by the Pauli Hamiltonian

$$(2) \quad H = \frac{1}{2m} [\boldsymbol{\pi}^2 - eB\sigma_3],$$

where $B = \text{rot } \mathbf{A} \equiv \epsilon^{ij} \partial_i A_j$ is the scalar magnetic field. It is now easy to see that the Hamiltonian is a perfect square in two different ways: both operators ⁽¹⁾

$$(3) \quad Q = \frac{1}{\sqrt{2m}} \boldsymbol{\pi} \cdot \boldsymbol{\sigma} \quad \text{and} \quad Q^* = \frac{1}{\sqrt{2m}} \boldsymbol{\pi} \times \boldsymbol{\sigma},$$

where $\boldsymbol{\sigma} = (\sigma_1, \sigma_2)$, satisfy

$$(4) \quad \{Q, Q\} = \{Q^*, Q^*\} = 2H.$$

Thus, for any static, purely magnetic field in the plane, H is an $N=2$ supersymmetric Hamiltonian. The supercharge Q is a standard object used in supersymmetric quantum mechanics [5]; the ‘twisted’ charge Q^* was used, e.g., by Jackiw [8], to describe the Landau states in a constant magnetic field — a classic example of supersymmetric quantum mechanics [5,7].

Let us assume henceforth that B is the field of a point-like magnetic vortex directed along the z -axis, $B = \Phi \delta(\mathbf{r})$, where Φ is the total magnetic flux. This can be viewed as an idealization of the spinning version of the Aharonov-Bohm experiment [9].

Inserting $A_i(\mathbf{r}) = -(\Phi/2\pi) \epsilon_{ij} \mathbf{r}^j / r^2$ into the Pauli Hamiltonian H , it is straightforward to check that $D = tH - \frac{1}{4} \{\boldsymbol{\pi}, \mathbf{r}\}$ and $K = -t^2 H + 2tD + \frac{1}{2} m r^2$ generate, along with H , the $\mathfrak{o}(2, 1)$ Lie algebra: $[D, H] = -iH$, $[D, K] = iK$, $[H, K] = 2iD$. The angular momentum, $J = \mathbf{r} \times \boldsymbol{\pi}$, adds to this $\mathfrak{o}(2, 1)$ an extra $\mathfrak{o}(2)$. (The correct definition of angular momentum requires boundary conditions, see [10]).

Commuting Q and Q^* with the expansion, K , yields two more generators, namely

$$(5) \quad S = i[Q, K] = \sqrt{\frac{m}{2}} \left(\mathbf{r} - \frac{\boldsymbol{\pi}}{m} t \right) \cdot \boldsymbol{\sigma}, \quad S^* = i[Q^*, K] = \sqrt{\frac{m}{2}} \left(\mathbf{r} - \frac{\boldsymbol{\pi}}{m} t \right) \times \boldsymbol{\sigma}.$$

It is now straightforward to see that both sets Q, S and Q^*, S^* extend the $\mathfrak{o}(2, 1) \cong \mathfrak{osp}(1/0)$ into an $\mathfrak{osp}(1/1)$ superalgebra. However, these two algebras do not close yet: the ‘mixed’ anticommutators $\{Q, S^*\}$ and $\{Q^*, S\}$ bring in a new conserved charge, *viz.* $\{Q, S^*\} = -\{Q^*, S\} = J + 2\Sigma$, where $\Sigma = \frac{1}{2} \sigma_3$. But J satisfies now non-trivial commutation relations with the supercharges, $[J, Q] = -iQ^*$, $[J, Q^*] = iQ$, $[J, S] = -iS^*$, $[J, S^*] = iS$. Thus, setting $Y = J + 2\Sigma = \mathbf{r} \times \boldsymbol{\pi} + \sigma_3$, the generators H, D, K, Y and Q, Q^*, S, S^*

⁽¹⁾ The cross product of two planar vectors, $\mathbf{u} \times \mathbf{v} = \epsilon_{ij} u^i v^j$, is a scalar.

satisfy

$$\begin{aligned}
 (6) \quad \begin{aligned}
 [Q, D] &= \frac{i}{2}Q, & [Q^*, D] &= \frac{i}{2}Q^*, \\
 [Q, K] &= -iS, & [Q^*, K] &= -iS^*, \\
 [Q, H] &= 0, & [Q^*, H] &= 0, \\
 [Q, Y] &= -iQ^*, & [Q^*, Y] &= iQ, \\
 [S, D] &= -\frac{i}{2}S, & [S^*, D] &= -\frac{i}{2}S^*, \\
 [S, K] &= 0, & [S^*, K] &= 0, \\
 [S, H] &= iQ, & [S^*, H] &= iQ^*, \\
 [S, Y] &= -iS^*, & [S^*, Y] &= iS, \\
 \{Q, Q\} &= 2H, & \{Q^*, Q^*\} &= 2H, \\
 \{S, S\} &= 2K, & \{S^*, S^*\} &= 2K, \\
 \{Q, Q^*\} &= 0, & \{S, S^*\} &= 0, \\
 \{Q, S\} &= -2D, & \{Q^*, S^*\} &= -2D, \\
 \{Q, S^*\} &= Y, & \{Q^*, S\} &= -Y.
 \end{aligned}
 \end{aligned}$$

When added to the $\mathfrak{o}(2, 1)$ relations, this means that our generators span the $\mathfrak{osp}(1/2)$ superalgebra [2]. On the other hand, $Z = J + \Sigma = \mathbf{r} \times \boldsymbol{\pi} + \frac{1}{2}\sigma_3$ commutes with all generators of $\mathfrak{osp}(1/2)$, so that the full symmetry is the direct product $\mathfrak{osp}(1/2) \times \mathfrak{o}(2)$, generated by

$$(7) \quad \begin{cases}
 Y = \mathbf{r} \times \boldsymbol{\pi} + \sigma_3, & Q = \frac{1}{\sqrt{2m}} \boldsymbol{\pi} \cdot \boldsymbol{\sigma}, \\
 H = \frac{1}{2m} [\boldsymbol{\pi}^2 - eB\sigma_3], & Q^* = \frac{1}{\sqrt{2m}} \boldsymbol{\pi} \times \boldsymbol{\sigma}, \\
 D = -\frac{1}{4} \{\boldsymbol{\pi}, \mathbf{q}\} - t \frac{eB}{2m} \sigma_3, & S = \sqrt{\frac{m}{2}} \mathbf{q} \cdot \boldsymbol{\sigma}, \\
 K = \frac{1}{2} m \mathbf{q}^2, & S^* = \sqrt{\frac{m}{2}} \mathbf{q} \times \boldsymbol{\sigma}, \\
 Z = \mathbf{r} \times \boldsymbol{\pi} + \frac{1}{2} \sigma_3,
 \end{cases}$$

where we have put $\mathbf{q} = \mathbf{r}(\boldsymbol{\pi}/m)t$.

The supersymmetric Hamiltonian (1) is the square of Jackiw's [8] two-dimensional Dirac operator $\boldsymbol{\pi} \times \boldsymbol{\sigma}$. But the Dirac operator is supersymmetric in any even dimensional space. The energy levels are therefore non-negative; eigenstates with non-zero energy are doubly degenerate; the system has $\text{Ent}(e\Phi - 1)$ zero-modes [8, 9]. The superalgebra (6) allows for a complete group-theoretical solution of the Pauli equation, along the lines indicated by D'Hoker and Vinet [2]. Details will be given elsewhere.

Notice that Jackiw's two-dimensional Dirac operator $\boldsymbol{\pi} \times \boldsymbol{\sigma}$ in Ref. [8] — essentially our Q^* — is associated with the unusual choice of the two-dimensional 'Dirac' (i.e. Pauli) matrices $\gamma_1^* = -\sigma_2$, $\gamma_2^* = \sigma_1$. Our helicity operator, Q , is again a 'Dirac operator' — but one associated with the standard choice $\gamma_1 = \sigma_1$, $\gamma_2 = \sigma_2$.

Note added. After this paper was completed and even submitted, we became aware of some papers [11] which expressed similar ideas. Our paper has consequently remained unpublished; parts of it entered [12].

References

- ¹ R. Jackiw, Ann. Phys. (N.Y.) **129**, 183 (1980).
- ² E. D'Hoker and L. Vinet, Phys. Lett. **137B**, 72 (1984); Comm. Math. Phys. **97**, 391 (1985).
- ³ R. Jackiw, Ann. Phys. (N.Y.) **201**, 83 (1990).
- ⁴ Y. Aharonov and D. Bohm, Phys. Rev. **115**, 485 (1959).
- ⁵ E. Witten, Nucl. Phys. **B185**, 513 (1981); P. Salomonson and J.W. Van Holten, Nucl. Phys. **B169**, 509 (1982); M. De Crombrugghe and V. Rittenberg, Ann. Phys. (N.Y.) **151**, 99 (1983).
- ⁶ M. Leblanc, G. Lozano and H. Min, Ann. Phys. (N.Y.) **219**, 328 (1992); C. Duval and P.A. Horváthy, submitted to Comm. Math. Phys. The bosonic galilean symmetry was noticed by R. Jackiw and S.-Y. Pi, Phys. Rev. **D 42**, 3500 (1990).
- ⁷ R. J. Hughes, V. A. Kostecký and M. M. Nieto, Phys. Rev. **D34**, 1100 (1986); E. D'Hoker, V. A. Kostecký and L. Vinet, in *Dynamical groups and spectrum generating algebras*, A. Bohm, Y. Ne'eman and A. O. Barut (eds), Vol. 1, p. 339; Singapore: World Scientific (1988).
- ⁸ R. Jackiw, Phys. Rev. **D29**, 2375 (1984).
- ⁹ C. R. Hagen, Phys. Rev. Lett. **64**, 503 (1990); R. Musto, L. O'Raifeartaigh and A. Wipf, Phys. Lett. **B 175**, 433 (1986); P. Forgács, L. O'Raifeartaigh and A. Wipf, Nucl. Phys. **B 293**, 559 (1987).
- ¹⁰ P.A. Horváthy, Phys. Rev. **A31**, 1151 (1985); F. Wilczek, Phys. Rev. Lett. **48**, 1144 (1982); R. Jackiw and A.N. Redlich, Phys. Rev. Lett. **50**, 555 (1983) W.C. Henneberger, Phys. Rev. Lett. **52**, 573 (1984).
- ¹¹ C. J. Park, Nucl. Phys. **376**, 99 (1992); J.-G. Demers, Mod. Phys. Lett. **8**, 827 (1993).
- ¹² C. Duval and P. A. Horváthy, J. Math. Phys. **35**, 2516 (1994) [hep-th/0508079]